

Appendix to the paper “Random walks on free products of cyclic groups”

Jean MAIRESSE ^{*} and Frédéric MATHÉUS [†]

February 2, 2008

Abstract

This paper is an appendix to the paper “Random walks on free products of cyclic groups” by J. Mairesse and F. Mathéus. It contains the details of the computations and the proofs of the results concerning the examples treated there.

Keywords: random walk, free product of finite groups, harmonic measure, drift, entropy, extremal generators.

AMS classification (2000): Primary 60J10, 60B15, 60J22, 65C40; Secondary 28D20; 37M25.

1 Introduction

Let G be a free product of a finite family of finite groups, and let the set of generators Σ be the union of the finite groups. We consider a transient nearest neighbor random walk on (G, Σ) . The harmonic measure is Markovian with a special combinatorial structure that can be completely described via a finite set of polynomial equations (see [1] for a proof and references). In several simple cases of interest, the polynomial equations can be explicitly solved, to get closed form formulas for the drift, the entropy, the probability of ever hitting an element, or the minimal positive harmonic functions of the walk. To illustrate, we give in [1] explicit formulas for the drift in the following cases: the modular group $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$, $\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$, $\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$ and the Hecke groups $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$. The present paper is an appendix to [1]. The purpose is to give the details of the computations and the proofs of the results concerning these examples. Vershik’s notion of extremal generators is also investigated in [1], and several examples and counter-examples are given. Again, the details of the computations are to be found in this appendix.

This appendix is self-contained in the following sense: the examples treated and the results proved are specified. On the other hand, we make a constant use of notations, notions or results from [1] which are not redefined or reproved. Hence, it is probably not a good idea to read this appendix independently of the parent paper.

2 Explicit Drift Computations

We give here the details of the computations and the complete proofs of the results stated in [1, §4.2-4.5].

^{*}LIAFA, CNRS-Université Paris 7, case 7014, 2, place Jussieu, 75251 Paris Cedex 05, France. E-mail: Jean.Mairesse@liafa.jussieu.fr

[†]LMAM, Université de Bretagne-Sud, Campus de Tohannic, BP 573, 56017 Vannes, France. E-mail: Frederic.Matheus@univ-ubs.fr

2.1 Random walks on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$

The group $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$ is isomorphic to the modular group $\text{PSL}(2, \mathbb{Z})$, i.e. the group of 2×2 matrices with integer entries and determinant 1, quotiented by $\pm \text{Id}$. Let a and b be the respective generators of $\mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}/3\mathbb{Z}$. A possible faithful representation of the group is

$$a = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}. \quad (1)$$

Consider a general nearest neighbor random walk $(\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}, \mu)$. Set $\mu(a) = 1 - p - q$, $\mu(b) = p$, $\mu(b^2) = q$. In Figure 1, we have represented the Cayley graph of $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$ and the one-step transitions of the random walk.




Figure 1: A nearest neighbor random walk on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$.

Consider first the case of a random walk satisfying: $\mu(a) = 1 - 2p$, $\mu(b) = \mu(b^2) = p$, $p \in (0, 1/2)$. Here, we are in the situation described at the beginning of [1, §4.1] where computing the drift is elementary and does not require any information on the harmonic measure. According to [1, Eq. (21)], we have:

$$\gamma = \frac{2p(1 - 2p)}{2 - p}. \quad (2)$$

To go further, let us compute explicitly the harmonic measure. Consider the Traffic Equations [1, Eq. (11)]. By symmetry, we should look for a solution r satisfying $r(b) = r(b^2)$. With this simplification, the Traffic Equations yield a single equation: $r(a) = 2(1 - p)r(b)$. So the solution is:

$$[r(a), r(b), r(b^2)] = \left[\frac{1 - p}{2 - p}, \frac{1}{2(2 - p)}, \frac{1}{2(2 - p)} \right].$$

Observe that the transition matrix defining the Markovian multiplicative harmonic measure is (the elements of Σ being ranked in the order $\{a, b, b^2\}$):

$$P = \begin{bmatrix} 0 & 1/2 & 1/2 \\ 1 & 0 & 0 \\ 1 & 0 & 0 \end{bmatrix}.$$

Hence the harmonic measure is the uniform measure except for the distribution of the initial element of the normal form. Observe that the harmonic measure is never stationary. Indeed,

the stationary distribution of P is $\pi = [1/2, 1/4, 1/4]$, and the equation $r = \pi$ has no solution in p .

Now let us consider the general nearest neighbor random walk on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$ defined by $\mu(a) = 1 - p - q, \mu(b) = p, \mu(b^2) = q$. The direct computation of the drift is not possible anymore. However, [1, Theorem 3.5] applies and, furthermore, the harmonic measure can be explicitly determined. Then we derive the drift using [1, Corollary 3.6].




Figure 2: The drift of the random walk $(\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}, \mu)$ as a function of $p = \mu(b)$ and $q = \mu(b^2)$, from two different angles.

The Traffic Equations [1, Eq. (11)] are:

$$\begin{aligned} r(a) &= (1 - p - q)(r(b) + r(b^2)) + qr(b) + pr(b^2) \\ r(b) &= pr(a) + qr(b^2) + (1 - p - q) \frac{r(a)r(b)}{r(b) + r(b^2)} \\ r(b^2) &= qr(a) + pr(b) + (1 - p - q) \frac{r(a)r(b^2)}{r(b) + r(b^2)}. \end{aligned} \tag{3}$$

Substituting $1 - r(b) - r(b^2)$ for $r(a)$ in the first equation, we get $r(b^2) = (1 - (2 - p)r(b))/(2 - q)$. Plugging this into the second (or third) equation allows us to compute $r(b)$. After some simplifications, we get:

$$\begin{aligned} r(a) &= \frac{p^2 + q^2 - 2pq - p - q + 4 - \Delta_1}{2\Delta_2} \\ r(b) &= \frac{q^3 - 3q^2 + p^2q - 5pq + 2p + 6q - (2 - q)\Delta_1}{2(q - p)\Delta_2} \\ r(b^2) &= \frac{p^3 - 3p^2 + pq^2 - 5pq + 6p + 2q - (2 - p)\Delta_1}{2(p - q)\Delta_2}, \end{aligned}$$

with

$$\begin{aligned} \Delta_1 &= \sqrt{p^4 + q^4 - 2p^3 - 2q^3 + 2p^2q^2 - 6p^2q - 6pq^2 + 5p^2 + 5q^2 + 6pq} \\ \Delta_2 &= p^2 + q^2 - pq - 2p - 2q + 4. \end{aligned}$$

Set $r = \mu(a) = 1 - p - q$. The drift is then:

$$\gamma = \frac{2r \left(pq - p - q + \sqrt{(p^2 + q^2)(3 + (r + p)^2 + (r + q)^2) + 2pq(2r + 1)} \right)}{(r + p)^2 + (r + q)^2 - pq + 2}. \quad (4)$$

For instance, the drift is maximized for $r = z_0, p = 1 - z_0, q = 0$ (or $r = z_0, p = 0, q = 1 - z_0$), where z_0 is the root of $[z^6 + 12z^4 - 4z^3 + 47z^2 - 48z + 12]$ whose numerical value is $0.490275 \dots$. The corresponding numerical value of the drift is $\gamma_{\max} = 0.163379 \dots$. This was not a priori obvious!

We are in the domain of application of [1, Proposition 3.9]. On the other hand, the harmonic measure is never shift-invariant. This can be proved by showing directly that the Stationary Traffic Equations have no solution.

2.2 Random walks on $\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$




Figure 3: A nearest neighbor random walk on $\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$.

Consider the probability μ such that $\mu(a) = \mu(b) = p, \mu(a^2) = \mu(b^2) = q = 1/2 - p$. According to [1, Proposition 3.8], the harmonic measure is ergodic. Let us solve the Stationary Traffic Equations [1, Eq. (12)]. By obvious symmetry, we should have $r(a) = r(b)$ and $r(a^2) = r(b^2)$. Set $r(i) = r(a^i) + r(b^i)$ for $i = 1, 2$. Using that $r(2) = 1 - r(1)$, the Stationary Traffic Equations yield a single equation:

$$(2p - \frac{1}{2})r(1)^2 + (2p - \frac{3}{2})r(1) - \frac{1}{2} = 0.$$

The unique solution lying between 0 and 1 gives

$$r(a) = r(b) = \frac{4p - 3 + \sqrt{16p^2 - 8p + 5}}{4(4p - 1)}, \quad r(a^2) = r(b^2) = \frac{4p + 1 - \sqrt{16p^2 - 8p + 5}}{4(4p - 1)}. \quad (5)$$

The harmonic measure is the ergodic Markovian multiplicative measure associated with r . In particular, the drift is

$$\gamma = \frac{1}{2} - p + (2p - \frac{1}{2})r(1) = -\frac{1}{4} + \frac{1}{4}\sqrt{16p^2 - 8p + 5}. \quad (6)$$

Now consider the nearest neighbour random walk $(\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}, \mu)$ with another symmetry: $\mu(a) = \mu(a^2) = p, \mu(b) = \mu(b^2) = q = 1/2 - p$. Here, it is not difficult to check that there is no solution to the Stationary Traffic Equations [1, Eq. (12)]. Let us turn our attention to the general Traffic Equations [1, Eq. (11)]. By obvious symmetry, we should look for a solution r satisfying $r(a) = r(a^2)$ and $r(b) = r(b^2)$. It implies that $r(\Sigma \setminus \Sigma_a) = r(b) + r(b^2) = 2r(b)$ and $r(\Sigma \setminus \Sigma_b) = r(a) + r(a^2) = 2r(a)$. The Traffic Equations get simplified as follows:

$$r(a) = pr(a) + (2p + q)r(b), \quad r(b) = qr(b) + (2q + p)r(a).$$

The solution to the above is:

$$r(a) = r(a^2) = \frac{1 + 2p}{6}, \quad r(b) = r(b^2) = \frac{1 - p}{3}.$$

The harmonic measure is the Markovian multiplicative measure associated with r . We have $\mu^\infty(u_1 u_2 \cdots u_k \Sigma^\mathbb{N}) = r(u_1)(1/2)^{k-1}$, i.e. μ^∞ is the uniform measure except for the distribution of the initial element of the normal form. Using [1, Eq. (19)], the drift is

$$\gamma = 4pq = 2p(1 - 2p). \quad (7)$$

The above formula for γ could have been obtained using [1, Eq. (21)] since μ is uniform on each of the two cyclic groups.

We have represented in Figure 4 the drift as a function of p for the two above cases: (i) $\mu(a) = \mu(b) = p, \mu(a^2) = \mu(b^2) = 1/2 - p$ (drift1); (ii) $\mu(a) = \mu(a^2) = p, \mu(b) = \mu(b^2) = 1/2 - p$ (drift2).




Figure 4: The drift of $(\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}, \mu)$ as a function of p .

At last, consider the case $\mu(a) = p, \mu(a^2) = q$, and $\mu(b) = \mu(b^2) = (1 - p - q)/2$. Solving explicitly the Traffic Equations is feasible but provides formula which are too lengthy to be reproduced here. However, for the drift, several simplifications occur, and we obtain the following formula:

$$\gamma = 2(1 - p - q) \sqrt{\frac{p^2 + q^2 + pq}{p^2 + q^2 - 2pq + 3}}.$$

The graph of γ as a function of p and q looks like the one in Figure 2.

For the general nearest neighbor random walk on $\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$, we did not succeed in solving completely the Traffic Equations.

2.3 The simple random walks on $\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$

We now consider the whole family of groups $\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$, $k \geq 3$. However, we have to compromise by considering only simple random walks (with respect to a minimal set of generators). In Figure 5, we show this simple random walk in the case $\mathbb{Z}/4\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$.

Here we obtain “semi-explicit” formulas: we define recursively a family of polynoms of one variable $(F_k)_k$, and the harmonic measure is expressed as a function of the unique solution in $(0, 1)$ of $F_k = 1$. For $k \geq 6$, we have no closed form formula for this unique root.

Consider the free product $G_1 \star G_2 = \mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$. Set $\Sigma_1 = G_1 \setminus \{1_{G_1}\} = \{a, \dots, a^{k-1}\}$, $\Sigma_2 = G_2 \setminus \{1_{G_2}\} = \{b, \dots, b^{k-1}\}$, and $\Sigma = \Sigma_1 \sqcup \Sigma_2$. Consider the simple random walk $(\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}, \mu)$ with $\mu(a) = \mu(b) = \mu(a^{-1}) = \mu(b^{-1}) = 1/4$.




Figure 5: The simple random walk on $\mathbb{Z}/4\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$.

According to [1, Proposition 3.8], the harmonic measure is ergodic. To determine it, we need to solve the simpler Stationary Traffic Equations [1, Eq. (12)]. Set $r(i) = r(a^i) + r(b^i) = 2r(a^i) = 2r(b^i)$. The Stationary Traffic Equations are:

$$r = r \left[\frac{1}{4} \begin{bmatrix} 0 & 1 & & & \\ 1 & 0 & 1 & & \\ & \ddots & \ddots & \ddots & \\ & & 1 & 0 & 1 \\ & & & 1 & 0 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} 1 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 1 \\ \vdots & \vdots & & \vdots & \vdots \\ 1 & 0 & \cdots & 0 & 1 \\ 1 & 0 & \cdots & 0 & 1 \end{bmatrix} + \frac{1}{4} \begin{bmatrix} r(1) & r(2) & \cdots & r(k-1) \\ 0 & \cdots & \cdots & 0 \\ \vdots & & & \vdots \\ 0 & \cdots & \cdots & 0 \\ r(1) & r(2) & \cdots & r(k-1) \end{bmatrix} \right]. \quad (8)$$

If $[r(1), r(2), \dots, r(k-1)]$ is a solution to (8), then it is clear that $[r(k-1), \dots, r(2), r(1)]$ is also a solution. By uniqueness, we deduce that $r(i) = r(k-i)$ for all i . In particular, we have $r(1) = r(k-1)$. For convenience, set $r(0) = r(k) = 1$. Then the Equations (8) rewrite as:

$$2(2 - r(1))r = [r(0) + r(2), r(1) + r(3), \dots, r(k-3) + r(k-1), r(k-2) + r(k)], \quad (9)$$

that is, for all $i \in \{1, k-1\}$,

$$r(i+1) + r(i-1) = 2(2 - r(1))r(i). \quad (10)$$

Consider the applications $F_n : [0, 1] \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, defined by

$$F_0(x) = 1, \quad F_1(x) = x, \quad \forall n \geq 2, \quad F_n(x) = 2(2 - x)F_{n-1}(x) - F_{n-2}(x). \quad (11)$$

To illustrate, here are the first values of F_i :

$$\begin{aligned}
F_2(x) &= -2x^2 + 4x - 1 \\
F_3(x) &= 4x^3 - 16x^2 + 17x - 4 \\
F_4(x) &= -8x^4 + 48x^3 - 96x^2 + 72x - 15 \\
F_5(x) &= 16x^5 - 128x^4 + 380x^3 - 512x^2 + 301x - 56 \\
F_6(x) &= -32x^6 + 320x^5 - 1264x^4 + 2496x^3 - 2554x^2 + 1244x - 209
\end{aligned}$$

By construction, the unique solution $r \in \mathring{\mathcal{B}}$ of the Traffic Equations (10) satisfies $r(i) = F_i(r(1))$ for all $i \in \{1, \dots, k-1\}$. Hence, it is enough to determine $r(1)$. For convenience, set $x_k = r(1)$. The equalities $r(i) = r(k-i)$ for all $i \in \{1, \dots, k-1\}$ rewrite as $F_i(x_k) = F_{k-i}(x_k)$ for all $i \in \{1, \dots, k-1\}$. Now, using (11) twice, we get

$$\begin{aligned}
F_k(x_k) &= 2(2-x_k)F_{k-1}(x_k) - F_{k-2}(x_k) \\
&= 2(2-x_k)F_1(x_k) - F_2(x_k) \\
&= 2(2-x_k)F_1(x_k) - [2(2-x_k)F_1(x_k) - F_0(x_k)] = F_0(x_k) = 1.
\end{aligned}$$

Next lemma shows that the equality $F_k(x_k) = 1$ is actually a characterization of x_k .

Lemma 2.1. *For $k \geq 3$, the equation $F_k(x) = 1$ has a unique solution x_k in $(0, 1)$.*

Proof. Notice first that:

$$F_k\left(\frac{1}{3}\right) = \frac{1}{3^k}, \quad F_k(1) = 1. \quad (12)$$

Now, let us prove that there exists a point $x^k \in (1/3, 1)$ such that F_k is strictly increasing on $[0, x^k)$ and strictly decreasing on $(x^k, 1]$. In view of (12), the proof of the lemma will then follow directly.

Consider the matrix

$$A = \begin{bmatrix} 2(2-x) & -1 \\ 1 & 0 \end{bmatrix}, \quad (13)$$

for $x \in [0, 1]$. Rewriting (11) in matrix form, we get,

$$\begin{bmatrix} F_k \\ F_{k-1} \end{bmatrix} = A \begin{bmatrix} F_{k-1} \\ F_{k-2} \end{bmatrix} = A^{k-1} \begin{bmatrix} F_1 \\ F_0 \end{bmatrix}. \quad (14)$$

The eigenvalues of A are

$$\lambda_1 = 2 - x + \sqrt{(1-x)(3-x)}, \quad \lambda_2 = 2 - x - \sqrt{(1-x)(3-x)}. \quad (15)$$

and $\lambda_1 \neq \lambda_2$ for $x \neq 1$. By diagonalizing, we get for $x \neq 1$,

$$A = \frac{1}{\lambda_1 - \lambda_2} \begin{bmatrix} \lambda_1 & \lambda_2 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix} \begin{bmatrix} 1 & -\lambda_2 \\ -1 & \lambda_1 \end{bmatrix}. \quad (16)$$

Using (14) and (16), we obtain

$$F_k(x) = \begin{bmatrix} \frac{1}{\lambda_1 - \lambda_2} & \frac{1}{\lambda_1 - \lambda_2} \end{bmatrix} \begin{bmatrix} \lambda_1^k & 0 \\ 0 & \lambda_2^k \end{bmatrix} \begin{bmatrix} x - \lambda_2 \\ \lambda_1 - x \end{bmatrix} = \frac{x - \lambda_2}{\lambda_1 - \lambda_2} \lambda_1^k + \frac{\lambda_1 - x}{\lambda_1 - \lambda_2} \lambda_2^k. \quad (17)$$

By differentiating (17), we get

$$F'_k = \frac{\lambda_1^k \left[(x - 3 + 2(1-x)^{1/2}(3-x)^{1/2})k + 2 \right] + \lambda_2^k \left[(-x + 3 + 2(1-x)^{1/2}(3-x)^{1/2})k + 2 \right]}{2(3-x)^{3/2}(1-x)^{1/2}}. \quad (18)$$

Set $f_1(x) = x - 3 + 2(1-x)^{1/2}(3-x)^{1/2}$ and $f_2(x) = -x + 3 + 2(1-x)^{1/2}(3-x)^{1/2}$. It is easily checked that f_1 and f_2 are strictly decreasing on $[0, 1]$. Since $f_2(1) = 2$, we have $f_2 > 0$ on $[0, 1]$. On the other hand $f_1(1/3) = 0$, hence we have $f_1 > 0$ on $[0, 1/3]$ and $f_1 < 0$ on $(1/3, 1]$. Since we have $\lambda_1 > 0$ and $\lambda_2 > 0$ for all $x \in [0, 1]$, it follows immediately that $F'_k > 0$ on $[0, 1/3]$ and has at most one zero in $[0, 1]$. Now by differentiating (11), we get that

$$F'_k(x) = 2(2-x)F'_{k-1}(x) - 2F_{k-1}(x) - F'_{k-2}(x).$$

In particular $F'_k(1) = 2F'_{k-1}(1) - 2 - F'_{k-2}(1)$. It follows easily that for $k \geq 3$, we have $F'_k(1) < 0$. Consequently, F'_k has exactly one zero in $[0, 1]$ which belongs to $(1/3, 1)$. $\square$

To illustrate Lemma 2.1, we have represented in Figure 6, the functions F_9 and F_{10} .




Figure 6: The functions $F_9(x)$ and $F_{10}(x)$.

Let us collect the above results. The harmonic measure of the simple random walk $(\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}, \mu)$ is the ergodic Markovian multiplicative measure associated with:

$$r = [x_k, F_2(x_k), F_3(x_k), \dots, F_2(x_k), x_k]$$

where x_k is the unique solution in $(0, 1)$ of the equation $F_k(x) = 1$.

Now let us compute the drift γ_k using the harmonic measure, see [1, Corollary 3.8]. We have

$$\gamma_k = \sum_{a \in \Sigma} \frac{1}{4} [-r(a^{-1}) + \sum_{b \in \Sigma \setminus \Sigma_a} r(b)] = 4 \frac{1}{4} [-\frac{x_k}{2} + \frac{1}{2}] = \frac{1-x_k}{2}.$$

Lemma 2.2. *The drift γ_k is a strictly increasing function of k and satisfies $\lim_k \gamma_k = 1/3$.*

Proof. Fix $x \in (1/3, 1)$. Recall that $F_n(x) > 0$ for all n (Lemma 2.1). Consider the expression (17). We have $\lambda_1 > 1$ and $0 < \lambda_2 < 1$. It follows easily that $\lim_n F_n(x) = \lim_n \lambda_1^n (x - \lambda_2) / (\lambda_1 - \lambda_2) = +\infty$.

Define $R_n = F_{n+1}(x) / F_n(x)$ for $n \in \mathbb{N}$. Using (11), we obtain

$$R_0 = x, \quad \forall n \geq 1, \quad R_n = 2(2-x) - \frac{1}{R_{n-1}}.$$

Assume that there exists some n such that $R_n \geq 1$. Then we have $R_{n+1} \geq 2(2-x) - 1 > 1$, since $x < 1$. Since $\lim_n F_n(x) = +\infty$, we deduce that there exists $n^x \in \mathbb{N} \setminus \{0\}$ such that $F_n(x)$ is a strictly decreasing function of n for $n < n^x$, and a strictly increasing function of n for $n > n^x$.

Now, let us prove that $x_{k+1} < x_k$. We have $x_k \in (1/3, 1)$, $F_1(x_k) = F_{k-1}(x_k)$. It implies that $n^{x_k} \leq k - 1$. In particular $F_{k+1}(x_k) > F_k(x_k) = 1$. In view of the behavior of F_{k+1} , as described in the proof of Lemma 2.1, it implies indeed that $x_{k+1} < x_k$. Therefore, γ_k is a strictly increasing function of k .

Consider $x \in (1/3, 1)$. We have seen that $\lim_n F_n(x) = \infty$. So there exists N_x such that $\forall n > N_x, F_n(x) > 1$, and therefore $\forall n > N_x, x_n < x$. It implies immediately that $\lim_n x_n = 1/3$ and $\lim_n \gamma_n = 1/3$. $\square$

To summarize, we have proved the theorem below.

Theorem 2.3. *Consider the group $\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$, the generators of the two cyclic groups being respectively a and b . Consider the simple random walk $(\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}, \mu)$ with $\mu(a) = \mu(a^{-1}) = \mu(b) = \mu(b^{-1}) = 1/4$. The harmonic measure is the Markovian multiplicative measure associated with r : $\forall i \in \{1, \dots, k-1\}$, $r(a^i) = r(b^i) = F_i(x_k)/2$, where F_i is defined in (11) and x_k is the unique solution in $(0, 1)$ of $F_k(x) = 1$. The drift is $\gamma_k = (1 - x_k)/2$. It is a strictly increasing function of k and $\lim_k \gamma_k = 1/3$.*

To illustrate, let us give the vector r for $\mathbb{Z}/4\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$:

$$[r(a), r(a^2), r(a^3)] = [r(b), r(b^2), r(b^3)] = \left[\frac{3 - \sqrt{5}}{8}, \frac{\sqrt{5}}{4} - \frac{1}{2}, \frac{3 - \sqrt{5}}{8} \right]. \quad (19)$$

and for $\mathbb{Z}/5\mathbb{Z} \star \mathbb{Z}/5\mathbb{Z}$:

$$[r(a), r(a^2), r(a^3), r(a^4)] = [r(b), r(b^2), r(b^3), r(b^4)] = \left[\frac{5 - \sqrt{13}}{8}, \frac{\sqrt{13} - 3}{8}, \frac{\sqrt{13} - 3}{8}, \frac{5 - \sqrt{13}}{8} \right].$$

Now, here is a table of the first values of γ_k , given either in closed form or numerically when no closed form could be found. Set $\mathbb{Z}_k = \mathbb{Z}/k\mathbb{Z}$.

	$\mathbb{Z}_3 \star \mathbb{Z}_3$	$\mathbb{Z}_4 \star \mathbb{Z}_4$	$\mathbb{Z}_5 \star \mathbb{Z}_5$	$\mathbb{Z}_6 \star \mathbb{Z}_6$	$\mathbb{Z}_7 \star \mathbb{Z}_7$	$\mathbb{Z}_8 \star \mathbb{Z}_8$
γ	1/4	$(\sqrt{5} - 1)/4$	$(\sqrt{13} - 1)/8$	0.330851...	0.332515...	0.333062...

The limit case. The limiting value of $1/3$ for γ_k as $k \rightarrow \infty$ was to be expected. Let us show why. The ‘limit’ of $\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$ as $k \rightarrow \infty$ is $\mathbb{Z} \star \mathbb{Z}$. The group $\mathbb{Z} \star \mathbb{Z}$ is the free group over two generators, its Cayley graph is $\mathbb{T}_4$, the homogeneous tree of degree 4. Analysing the simple random walk on $\mathbb{Z} \star \mathbb{Z}$ is straightforward. Let a and b be the respective generators of the two groups and let μ^∞ be the harmonic measure with respect to the set of generators $\Sigma = \{a, a^{-1}, b, b^{-1}\}$. (The existence of μ^∞ is classical, see the discussion following [1, Proposition 2.1].) A word $u \in \Sigma^*$ is a normal form (or reduced) word if $u = u_1 \cdots u_k$ with $\forall i, u_{i+1} \neq u_i^{-1}$. For any normal form word $u \in \Sigma^k$, we have, by symmetry, $\mu^\infty(u\Sigma^\mathbb{N}) = 1/4 \times 1/3^{k-1}$. In particular the drift is $\gamma = 1/2$.

However, our computations in $\mathbb{Z}/k\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$ correspond to the set of generators $\{a, \dots, a^{k-1}, b, \dots, b^{k-1}\}$, and not to the set of generators $\{a, a^{-1}, b, b^{-1}\}$. To get the analog in $\mathbb{Z} \star \mathbb{Z}$, we have to compute the drift with respect to the infinite set of generators $S = \{a^i, b^i, i \in \mathbb{Z} \setminus \{0\}\}$. Let $X_\infty = x_1 \cdots x_n \cdots$ be a r.v. distributed according to μ^∞ . Let $\tau = \max\{k \mid x_1 = \cdots = x_k\}$. The expectation $E[\tau]$ is the rate of reduction of the length when recoding a reduced word over Σ in terms of S . Therefore the drift with respect to S is equal to $(1/2 \times 1/E[\tau])$. Using the above formula for μ^∞ , we get $P\{\tau = k\} = 1/3^{k-1} \times 2/3$. It implies that $E[\tau] = 3/2$. So the drift of the simple random walk on $\mathbb{Z} \star \mathbb{Z}$ with respect to S is $(1/3 = 1/2 \times 2/3)$.

2.4 The simple random walks on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$

We study simple (for a minimal set of generators) random walks on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$. The results are “semi-explicit” as in §2.3.

The groups $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$ are known as the Hecke groups. A possible faithful representation of $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$, generalizing the one given in (1), is

$$a = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} 2\cos(\pi/k) & -1 \\ 1 & 0 \end{bmatrix},$$

where a and b are the generators of $\mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}/k\mathbb{Z}$ respectively.

Consider the free product $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$, with a and b the respective generators of the two cyclic groups. We consider the simple random walk $(\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}, \mu)$ with $\mu(a) = \mu(b) = \mu(b^{-1}) = 1/3$.




Figure 7: The simple random walk on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$.

We have seen above that even for $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/3\mathbb{Z}$, the harmonic measure is not ergodic. Hence we should try to solve the general Traffic Equations [1, Eq. (11)]:

$$\begin{aligned} r(a) &= \frac{1}{3} \left[1 - r(a) + r(b) + r(b^{k-1}) \right] \\ r(b) &= \frac{1}{3} \left[r(a) + r(b^2) + r(b) \frac{r(a)}{1 - r(a)} \right] \\ r(b^i) &= \frac{1}{3} \left[r(b^{i-1}) + r(b^{i+1}) + r(b^i) \frac{r(a)}{1 - r(a)} \right], \quad \forall i = 2, \dots, k-2 \\ r(b^{k-1}) &= \frac{1}{3} \left[r(b^{k-2}) + r(a) + r(b^{k-1}) \frac{r(a)}{1 - r(a)} \right]. \end{aligned}$$

By symmetry, we must have $r(b^i) = r(b^{k-i})$ for all i . Hence the first equation gives: $r(a) = 1/4 + r(b)/2$. Now set $r(b^0) = r(b^k) = r(a) = 1/4 + r(b)/2$. Rewriting the Traffic Equations (except the first one), we get for $i = 1, \dots, k-1$,

$$\left[\frac{3 - 2r(b)}{8(1 - r(b))} \right] r(b^i) = r(b^{i-1}) + r(b^{i+1}). \quad (20)$$

This should be compared with (10). Consider the applications $G_n : [0, 1] \rightarrow \mathbb{R}$, $n \in \mathbb{N}$, defined by

$$G_0(x) = \frac{1}{4} + \frac{x}{2}, \quad G_1(x) = x, \quad \forall n \geq 2, \quad G_n(x) = \frac{8(1-x)}{3-2x} G_{n-1}(x) - G_{n-2}(x). \quad (21)$$

In illustration, here are the first few values of G_i :

$$\begin{aligned}(2x-3)G_2(x) &= 7x^2 - 7x + 3/4 \\ (2x-3)^2G_3(x) &= 52x^3 - 100x^2 + 53x - 6 \\ (2x-3)^3G_4(x) &= 1552x^4 - 4416x^3 + 4296x^2 - 1600x + 165\end{aligned}$$

From now on, the arguments become very close to the ones in §2.3. Hence, we choose not to detail them. Observe just that the value of x playing a special role (similar to the one of $1/3$ in §2.3) is now $1/6$ with $G_i(1/6) = 1/(3 \times 2^i)$.

Next theorem summarizes the results.

Theorem 2.4. *Consider the group $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$, the generators of the two cyclic groups being respectively a and b . Consider the simple random walk $(\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}, \mu)$ with $\mu(a) = \mu(b) = \mu(b^{-1}) = 1/3$. Let the functions G_i be defined as in (21) and let y_k be the unique solution in $(0, 1/2)$ of $G_{k-1}(y_k) = y_k$. The harmonic measure is the Markovian multiplicative measure associated with r : $r(a) = G_0(y_k)$, $\forall i \in \{1, \dots, k-1\}$, $r(b^i) = G_i(y_k)$. The drift is $\gamma_k = (1 - 2y_k)/3$. It is a strictly increasing function of k and $\lim_k \gamma_k = 2/9$.*

Here is the vector r for $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$:

$$[r(a), r(b), r(b^2), r(b^3)] = \left[\frac{7-\sqrt{7}}{12}, \frac{2}{3} - \frac{\sqrt{7}}{6}, \frac{-11+5\sqrt{7}}{12}, \frac{2}{3} - \frac{\sqrt{7}}{6} \right].$$

and for $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/5\mathbb{Z}$:

$$[r(a), r(b), r(b^2), r(b^3), r(b^4)] = \left[\frac{21-\sqrt{61}}{38}, \frac{23}{38} - \frac{\sqrt{61}}{19}, \frac{-29+5\sqrt{61}}{76}, \frac{-29+5\sqrt{61}}{76}, \frac{23}{38} - \frac{\sqrt{61}}{19} \right].$$

In next table, we give the drift of the simple random walk on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$ for the small values of k . The drift is given in closed form when possible, otherwise numerically. Set $\mathbb{Z}_k = \mathbb{Z}/k\mathbb{Z}$.

	$\mathbb{Z}_2 \star \mathbb{Z}_3$	$\mathbb{Z}_2 \star \mathbb{Z}_4$	$\mathbb{Z}_2 \star \mathbb{Z}_5$	$\mathbb{Z}_2 \star \mathbb{Z}_6$	$\mathbb{Z}_2 \star \mathbb{Z}_7$	$\mathbb{Z}_2 \star \mathbb{Z}_8$
γ	$2/15$	$(\sqrt{7}-1)/9$	$(2\sqrt{61}-4)/57$	$0.213412\dots$	$0.217921\dots$	$0.220101\dots$

The limit case. As in §2.3, the result $\lim_k \gamma_k = 2/9$ can be recovered using only elementary arguments. The ‘limit’ as $k \rightarrow \infty$ of $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/k\mathbb{Z}$ is $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}$. Let a and b be the respective generators of $\mathbb{Z}/2\mathbb{Z}$ and $\mathbb{Z}$. The Cayley graph of $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}$ with respect to $\Sigma = \{a, b, b^{-1}\}$ is $\mathbb{T}_3$, the homogeneous tree of degree 3.

From there, we proceed as in §2.3. The drift of the simple random walk on $\mathbb{T}_3$ is obviously $1/3$. The harmonic measure is $\mu^\infty(u_1 \cdots u_k \Sigma^\mathbb{N}) = 1/3 \times 1/2^{k-1}$ for any normal form word $u_1 \cdots u_k \in \Sigma^k$. Set $S = \{a, b^i, i \in \mathbb{Z} \setminus \{0\}\}$. Let $X_\infty = x_1 \cdots x_n \cdots$ be a r.v. distributed according to μ^∞ . Let $\tau = \max\{k \mid x_1 = \cdots = x_k\}$. If $x_1 = a$, then $\tau = 1$. On the other hand, $P\{\tau = k \mid x_1 = b \text{ or } b^{-1}\} = 1/2^k$. So we have $E[\tau \mid x_1 = a] = 1$ and $E[\tau \mid x_1 \in \{b, b^{-1}\}] = 2$. In a normal form word over Σ , we have a strict alternance of a single a with blocks of b or b^{-1} . Hence the expected reduction in length when recoding a normal form word over S is $3/2$. We conclude that the drift of the simple random walk on $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}$ with respect to S is $(2/9 = 1/3 \times 2/3)$.

3 Entropy and Extremal Generators

Here, we give the proofs of the results announced in [1, §5.1], and recalled below.

Statement of the results.

• A. Consider a free product of two finite groups, $G_1 \star G_2$. Consider the probability μ_p on Σ such that $\mu_p = pU_{\Sigma_1} + (1-p)U_{\Sigma_2}$, where U_{Σ_i} is the uniform distribution on Σ_i , and where $p \in (0, 1)$. Then we have $h/(\gamma v) = 1$, for the whole family of probabilities μ_p . This does not extend to the free product of more than 2 groups, see B, nor to minimal sets of generators $S \subset \Sigma$, see C.

• B. Consider the group $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/2\mathbb{Z}$, and let a, b , and c be the non-identity elements of the three cyclic groups. Consider the family of probability measures $\mu_p, p \in (0, 1/2)$, defined by: $\mu_p(a) = \mu_p(b) = p, \mu_p(c) = 1 - 2p$. Among them, the only probability such that $h/(\gamma v) = 1$ is $\mu_{1/3}$.

• C. Consider the group $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$. Let a and b be respective generators of the two cyclic groups. The minimal set of generators $S = \{a, b, b^{-1}\}$ is extremal but $h/(\gamma v) < 1$ for all μ in $\mathcal{S}$. Consider now the group $\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$. The minimal set of generators $S = \{a, a^{-1}, b, b^{-1}\}$ is extremal. Actually, the only symmetric probability measure on S for which $h = \gamma v$ is $\mu = p(\delta_a + \delta_{a^{-1}}) + (1/2 - p)(\delta_b + \delta_{b^{-1}})$ with $p = 0.432692 \dots$ is the middle root of the polynomial $5x^3 - 13x^2 + 7x - 1$. Observe that $\mu \in \mathcal{S}$. Last, consider the group $\mathbb{Z}/4\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$. The minimal set of generators $S = \{a, a^{-1}, b, b^{-1}\}$ is not extremal. Indeed,

$$Q(S) = \frac{5 + \sqrt{5} \log(1/2 + \sqrt{5}/2)}{4 \log(1 + \sqrt{2})} = 0.987686 \dots$$

Proof of A. The volume is $v = \log(|\Sigma_1||\Sigma_2|)/2$. Set $r(\Sigma_i) = \sum_{x \in \Sigma_i} r(x)$ and $q(\Sigma_i) = \sum_{x \in \Sigma_i} q(x)$. By symmetry, for all $x \in \Sigma_i$, we have $r(x) = r(\Sigma_i)/|\Sigma_i|$ and $q(x) = q(\Sigma_i)/|\Sigma_i|$. Using [1, Eq. (7)] and [1, Eq. (36)] we obtain

$$\begin{aligned} \gamma &= \mu(\Sigma_1) \left(r(\Sigma_2) - \frac{r(\Sigma_1)}{|\Sigma_1|} \right) + \mu(\Sigma_2) \left(r(\Sigma_1) - \frac{r(\Sigma_2)}{|\Sigma_2|} \right) \\ h &= -\mu(\Sigma_1) \log\left(\frac{|\Sigma_1|}{q(\Sigma_1)}\right) \left(r(\Sigma_2) - \frac{r(\Sigma_1)}{|\Sigma_1|} \right) - \mu(\Sigma_2) \log\left(\frac{|\Sigma_2|}{q(\Sigma_2)}\right) \left(r(\Sigma_1) - \frac{r(\Sigma_2)}{|\Sigma_2|} \right). \end{aligned}$$

To go further, let us solve the Traffic Equations. We have:

$$r(\Sigma_1) = \frac{\mu(\Sigma_1) + \mu(\Sigma_2)/|\Sigma_2|}{1 + \mu(\Sigma_1)/|\Sigma_1| + \mu(\Sigma_2)/|\Sigma_2|}, \quad r(\Sigma_2) = \frac{\mu(\Sigma_2) + \mu(\Sigma_1)/|\Sigma_1|}{1 + \mu(\Sigma_1)/|\Sigma_1| + \mu(\Sigma_2)/|\Sigma_2|}.$$

It follows immediately that:

$$\begin{aligned} r(\Sigma_1) (\mu(\Sigma_2) + \mu(\Sigma_1)/|\Sigma_1|) &= r(\Sigma_2) (\mu(\Sigma_1) + \mu(\Sigma_2)/|\Sigma_2|) \\ \iff \mu(\Sigma_1) (r(\Sigma_2) - r(\Sigma_1)/|\Sigma_1|) &= \mu(\Sigma_2) (r(\Sigma_1) - r(\Sigma_2)/|\Sigma_2|). \end{aligned}$$

Using the above equality, we can rewrite the drift and the entropy as follows:

$$\begin{aligned} \gamma &= 2\mu(\Sigma_1) (r(\Sigma_2) - r(\Sigma_1)/|\Sigma_1|) \\ h &= -\mu(\Sigma_1) (r(\Sigma_2) - r(\Sigma_1)/|\Sigma_1|) (\log(|\Sigma_1|/q(\Sigma_1)) + \log(|\Sigma_2|/q(\Sigma_2))) \\ &= -\mu(\Sigma_1) (r(\Sigma_2) - r(\Sigma_1)/|\Sigma_1|) \log(|\Sigma_1||\Sigma_2|), \end{aligned}$$

where we have used the identity $q(\Sigma_1)q(\Sigma_2) = 1$ to obtain the last equality. We conclude that: $h/\gamma = \log(|\Sigma_1||\Sigma_2|)/2 = v$.

Proof of B. The volume of $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/2\mathbb{Z}$ is $v = \log(2)$. The Traffic Equations can be explicitly solved and we obtain the drift γ_p and the entropy h_p associated with the measure μ_p :

$$\begin{aligned}\gamma_p &= 3p^2 + \frac{3-9p}{2} + \frac{3p-1}{2} \sqrt{4p^2 - 20p + 9} \\ h_p &= \gamma_p \log(2) + \log(3 - 2p + \sqrt{4p^2 - 20p + 9}) \frac{p}{2} (-1 + 2p + \sqrt{4p^2 - 20p + 9}) + \\ &\quad \log\left(-1 + \frac{\sqrt{4p^2 - 20p + 9}}{1 - 2p}\right) \frac{1 - 2p}{2} (3 - 2p - \sqrt{4p^2 - 20p + 9}).\end{aligned}$$

Let us write $h_p = \gamma_p \log(2) + f(p)$. A direct analysis shows that $f(p) = 0$ if and only if $p = 1/3$. We deduce that $h_p/\gamma_p = v$ if and only if $p = 1/3$.

Proof of C. Consider first the group $\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$ and let a and b be the respective generators of the cyclic groups. Consider the minimal set of generators $S = \{a, b, b^{-1}\}$. First of all, let us compute the volume $v(S)$ with respect to S . Deciding to write b^2 as bb , we enumerate the group elements using the automaton of Figure 8. More precisely, each path of length n (starting with a , b , or b^{-1}) corresponds to a different group element of length $n + 1$ with respect to S . It




Figure 8: An automaton enumerating the group elements according to $|\cdot|_S$.

follows that $v(S) = \log(\rho)$ where ρ is the spectral radius of A , the matrix associated with the automaton. We deduce that $v(S) = \log(1/2 + \sqrt{5}/2)$.

Consider $\mu \in \mathcal{S}$ with $\mu(a) = 1 - 2p$, $\mu(b) = \mu(b^{-1}) = p$, $p \in (0, 1/2)$. The solution to the Traffic Equations is:

$$\begin{aligned}r(a) &= \frac{2 - p - p^2 - p\sqrt{5 - 6p + p^2}}{4(1 - p)} \\ r(b) = r(b^{-1}) &= \frac{3 - p - \sqrt{5 - 6p + p^2}}{4} \\ r(b^2) &= \frac{-4 + 5p - p^2 + (2 - p)\sqrt{5 - 6p + p^2}}{4(1 - p)}.\end{aligned}$$

The value of the drift follows (when computing the drift, one must remember that the element b^2 has length 2 with respect to S):

$$\gamma(S, \mu) = \frac{p(2p + 5)}{2} - \frac{p(2p + 1)\sqrt{5 - 6p + p^2}}{2(1 - p)}.$$

Knowing r , the explicit formula for $h(\mu)$ is computable, but quite long. One checks directly on the resulting expression that $h(\mu)/(\gamma(S, \mu)v(S)) < 1$ for all $p \in (0, 1/2)$, and that




Figure 9: The ratio $h(\mu)/(\gamma(S, \mu)v(S))$ for $(\mathbb{Z}/2\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}, \mu)$ (top) and $(\mathbb{Z}/4\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}, \mu)$ (bottom) as a function of $p = \mu(b) = \mu(b^{-1})$.

$\lim_{p \searrow 0} h(\mu)/(\gamma(S, \mu)v(S)) = 1$. See Figure 9. Hence $Q(S) = 1$, the set of generators S is extremal.

Consider now the group $\mathbb{Z}/3\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$ and the minimal set of generators $S = \{a, a^{-1}, b, b^{-1}\}$. With the same type of arguments as above, we obtain that the volume with respect to S is $v(S) = \log(\rho)$ where $\rho = 2.21432 \dots$ is the real root of the polynomial $x^3 - 4x - 2$.

Consider the measure $\mu \in \mathcal{S}$ defined by $\mu(a) = \mu(a^{-1}) = p, \mu(b) = \mu(b^{-1}) = 1/2 - p, p \in (0, 1/2)$. The solution to the Traffic Equations is:

$$\begin{aligned} r(a) = r(a^{-1}) &= \frac{5 + 4p - 4p^2 - (1 - 2p)\sqrt{9 + 4p - 4p^2}}{8(2 - p)(1 + p)} \\ r(b) = r(b^{-1}) &= \frac{5 - 2p - \sqrt{9 + 4p - 4p^2}}{4(2 - p)} \\ r(b^2) &= \frac{-7 - 6p + 4p^2 + 3\sqrt{9 + 4p - 4p^2}}{4(2 - p)(1 + p)}. \end{aligned}$$

Computing the drift cautiously as above gives:

$$\gamma(S, \mu) = \frac{(1 - 2p)(12 + 11p - 10p^2 - (4 - 5p)\sqrt{9 + 4p - 4p^2})}{4(2 - p)(1 + p)}.$$

With the explicit formula for $h(\mu)$, one checks directly that the maximum of $h(\mu)/(\gamma(S, \mu)v(S))$ is 1 and is obtained for $p = 0.432692 \dots$ which is the middle root of the polynomial $5x^3 - 13x^2 + 7x - 1$.

Last, consider the group $\mathbb{Z}/4\mathbb{Z} \star \mathbb{Z}/4\mathbb{Z}$ and the minimal set of generators $S = \{a, a^{-1}, b, b^{-1}\}$. Again, with the same type of arguments as above, we obtain that the volume with respect to S is $v(S) = \log(1 + \sqrt{2})$.

Consider the measure defined by $\mu(a) = \mu(a^{-1}) = p, \mu(b) = \mu(b^{-1}) = 1/2 - p$ for some $p \in (0, 1/2)$. The solution to the Traffic Equations is obtained formally. Consider the two equations:

$$\begin{aligned} (-4 + 8p^2)X^4 + (4 + 8p^2)X^3 + (3 + 4p)X^2 - (2 + 2p)X - (1 + 2p) &= 0 \\ (-2 - 8p + 8p^2)X^4 + (6 - 8p + 8p^2)X^3 + (5 - 4p)X^2 + (-3 + 2p)X + 2p - 2 &= 0 \end{aligned}$$

Denote respectively by T and U the largest real root of the two equations. (The exact formulas for T and U are too long to be reproduced here.) We have:

$$r(a) = r(a^{-1}) = \frac{1}{2(1+U)}, \quad r(a^2) = \frac{1}{2U(1+U)}, \quad r(b) = r(b^{-1}) = \frac{1}{2(1+T)}, \quad r(b^2) = \frac{1}{2T(1+T)}.$$

The value of the drift and the entropy follows readily. We get:

$$\begin{aligned} \gamma(S, \mu) &= \frac{1 - 4p + 2U(1 - 2p)}{2U(1 + U)} + \frac{-1 + 4p + 4pT}{2T(1 + T)} \\ h(\mu) &= \log\left(\frac{(U+1)(2T+1)}{T(T+1)}\right) \frac{p(2U^2 - U - 1)}{U(U+1)} + \log(U) \frac{p(U-1)}{U(U+1)} \\ &\quad + \log\left(\frac{(T+1)(2U+1)}{U(U+1)}\right) \frac{(1/2 - p)(2T^2 - T - 1)}{T(T+1)} + \log(T) \frac{(1/2 - p)(T-1)}{T(T+1)}. \end{aligned}$$

The ratio $h(\mu)/(\gamma(S, \mu)v(S))$ as a function of p has been represented in Figure 9. It is maximized for $p = 1/4$. This was to be expected since the two groups in the free product are isomorphic. For $p = 1/4$, one has: $r(a) = r(b) = r(a^{-1}) = r(b^{-1}) = (3 - \sqrt{5})/4$, $r(a^2) = r(b^2) = -1 + \sqrt{5}/2$, see (19). Hence, $\gamma(S, \mu) = (3 - \sqrt{5})/2$ and $h(\mu) = (5 - \sqrt{5}) \log(1/2 + \sqrt{5}/2)/4$. It follows that:

$$Q(S) = \frac{5 + \sqrt{5}}{4} \frac{\log(1/2 + \sqrt{5}/2)}{\log(1 + \sqrt{2})} = 0.987686 \dots$$

In particular, the generators S are not extremal.

References

- [1] J. Mairesse and F. Mathéus. Random walks on free products of cyclic groups. arXiv:math.PR/0509211, 2005.